

Diatom 3: 3-8 (英文)

根来健一郎：日本の硅藻学への貢献 (論文目録)

Prof. Dr. Ken-ichiro Negoro : Contributions to the Diatomology of Japan (Bibliography)

- 1) Negoro, K. (1937) : Die Algen aus dem Kamagaike, Hochmoorteich, in Nagano Prefäktur. Jap. Jour. Limnol., Vol. **7**, pp. 121-126. (In Japanese, with German summary).
- 2) Negoro, K. (1938) : Die Algen aus dem *Sphagnum*-Sumpf von Usagizima bei Nikko. Jap. Jour. Limnol., Vol. **8**, pp. 9-13. (In Japanese, with German summary).
- 3) Negoro, K. (1938) : A species of diatom living in the bottom mud of Lake Kata-numa, a very strongly acid-water lake in Japan. Bot. & Zool., Vol. **6**, pp. 1731-1736. (In Japanese).
- 4) Negoro, K. (1940) : The diatom-flora of the Nasu Hot Springs (Preliminary report). Bot. Mag. (Tokyo), Vol. **54**, pp. 63-65. (In Japanese).
- 5) Negoro, K. (1940) : Some diatoms from Itibisinai-ko, a mineralogenous acidotrophic lake in Kunasiri-zima of the South Kurile Islands. Bot. Mag. (Tokyo), Vol. **54**, pp. 106-109. (In Japanese, with English summary).
- 6) Negoro, K. (1940) : A *Pinnularia*-diatom from Okama, the acidotrophic crater lake of Volcano Zaô. Jap. Jour. Limnol., Vol. **9**, pp. 176-179. (In Japanese).
- 7) Negoro, K. (1941) : Über die allgemeine Verbreitung und das massenhafte Vorkommen von *Pinnularia Brauni* var. *amphicephala* (A. Mayer) Cleve in den mineralogen-azidotrophen Gewässern Japans. Proc. Imp. Acad. Tokyo, Vol. **17**, pp. 425-428.
- 8) Negoro, K. (1942) : The habitats and the circumstances of *Pinnularia acoricola* Hustedt in Japan. Med. & Biol., Vol. **1**, pp. 111-114. (In Japanese).
- 9) Negoro, K. (1942) : Diatoms from the Shimogamo Hot Springs, Shizuoka Prefecture. Med. & Biol., Vol. **1**, pp. 211-215. (In Japanese).
- 10) Negoro, K. (1942) : Die Diatomeenvegetation der Isobe-Mineralquellen. Bot. Mag. (Tokyo), Vol. **56**, pp. 392-404. (In Japanese, with German summary).
- 11) Negoro, K. (1943) : Über die Algenvegetation der Thermen von Kusatu, Gunma Präfektur. Bot. Mag. (Tokyo), Vol. **67**, pp. 302-312. (In Japanese, with German summary).
- 12) Negoro, K. (1944) : Untersuchungen über die vegetation der mineralogen-azidotrophen gewässer Japans. Sci Rep. Tokyo Bunrika Daigaku, Sec. B, Vol. **6**, No. 101, pp. 231-374.
- 13) Negoro, K. (1949) : The diatom-vegetation in the hydrogen sulphidehot-springs at Yumoto, Nikko. Jap. Jour. Limnol., Vol. **14**, pp. 15-21. (In Japanese)
- 14) Negoro, K. (1952) : The phytoplankton of the Fukami Pond, Nagano Prefecture. M. Uéno's "The Fukami Pond", pp. 39-42. (In Japanese).
- 15) Negoro, K. (1952) : Diatoms from the Nitta Pond, Nagano Prefecture. M. Uéno's "The Fukami Pond", pp. 87-91. (In Japanese).
- 16) Negoro, K. (1953) : Preliminary report on the diatom-flora of the Ozé High Moors. Sôrui

- (Bull. Jap. Soc. Phycol.), Vol. **1**, pp. 41-44. (In Japanese).
- 17) Negoro, K. (1954) : The Algae from paddy-fields of Province Kii. I. Diatom-flora of the paddy-fields at Kemi, Wakayama City. Nanki-Seibutsu, Vol. **5**, pp. 1-4. (In Japanese).
 - 18) Negoro, K. (1954) : Two noticeable diatoms from Lake Biwa. Kagaku, Vol. **24**, p. 420. (In Japanese).
 - 19) Negoro, K. (1954) : Diatom Shells in the bottom deposits of the profound zone of Lake Biwa. Kagaku, Vol. **24**, pp. 527-528. (In Japanese).
 - 20) Negoro, K. (1954) : The Phytoplankton of the Matsubara Lake Groups, Nagano Prefecture. M. Uéno's "The Matsubara Lake Groups", pp. 327-331. (In Japanese).
 - 21) Negoro, K. (1954) : The Plankton of Lake Biwa. Sci. Rep. Fish. Exp. Stat. Shiga Pref., Spec. Publ., pp. 1-40, 10 pls. with 101 figs. (In Japanese).
 - 22) Negoro, K. (1955) : Diatom flora of the Pond Sokonashi of the Tokara Islands (A preliminary report). Jap. Jour. Limnol., Vol. **17**, pp.49-52. (In Japanese, with English summary).
 - 23) Negoro, K. (1956) : The Phytoplankton of the main basin of Lake Biwa. Jap. Jour. Limnol., Vol. **18**, pp. 37-46. (In Japanese, with English summary).
 - 24) Negoro, K. (1956) : The diatom shells in the bottom deposits of Lake Yogo, north of Lake Biwa. Jap. Jour. Limnol., Vol. **18**, pp. 134-140. (In Japanese, with English summary).
 - 25) Negoro, K. (1957) : The plankton of Lake Ibanaike (or Nakanoumi) as trophic indicators. Jap. Jour. Limnol., Vol. **19**, pp. 68-71. (In Japanese, with English summary).
 - 26) Negoro, K. (1957) : The algal vegetation of the hot springs of Asama, Hongô-mura, Nagano Prefecture. History and Nature of the Hongô-Village, Vol. **4**, pp. 63-64. (In Japanese).
 - 27) Negoro, K. (1958) : An analytical study of diatom shells in the bottom deposits of Lake Biwa-ko. Jap. Jour. Limnol., Vol. **19**, pp. 77-84. (In Japanese, with English summary).
 - 28) Negoro, K. (1958) : Diatom shells in the bottom deposits of some lakes of Shimominochi District, Nagano Prefecture. M. Uéno's "The Lakes of Shimominochi District, Nagano Prefecture. M. Uéno's "The Lakes of Shimominochi, North Shinano, Middle Japan", pp. 157-166. (In Japanese).
 - 29) Negoro, K. (1959) : Some noticeable plankton-algae of Lake Biwa and its effluent. Algal indicators of a water system. Mem. Coll. Sci. Univ. Kyoto, Ser. B, Vol. **26**, No.3, pp. 311-314.
 - 30) Negoro, K. (1959) : The chemical quality of water and the diatom flora of Lake Otori-ike and its neighbouring waters. Jap. Jour. Limnol., Vol. **20**, pp. 151-159. (In Japanese, with English summary).
 - 31) Negoro, K. (1960) : Studies on the diatom-vegetation of Lake Biwa-ko (First report). Jap.

- Jour. Limnol., Vol. **21**, pp. 200-220. (In Japanese, with English summary).
- 32) Negoro, K. (1961) : Diatoms from some inland-waters of Antarctica (Preliminary report). The Antarctic Record, No. **11**, pp. 872-873.
- 33) Negoro, K. (1962) : Algae and mosses in the mineralogenous acidotrophic inland-waters of Japan. Acta Phytotax. Geobot., Vol. **20**, pp. 314-321. (In Japanese).
- 34) Negoro, K. (1964) : On the stratum of marine origin in the bottom deposits of Lake Kasumi-ga-ura. An analytical study of microfossils. Physiol. & Ecol., Vol. **12**, pp. 191-198. (In Japanese, with English summary).
- 35) Negoro, K. (1967) : First report of the regular limnological survey of Lake Biwa (Oct. 1965-Dec. 1966). □. Plankton in general and phytoplankton. Mem. Fac. Sci. Kyoto Univ., Ser. Biol., Vol. I, No. 1, pp. 41-61.
- 36) Negoro, K. (1967) : An analytical study of diatom shells in the bottom deposits of Lake Biwa-ko, based on a new core-sample. Jap. Jour. Limnol., Vol. **28**, pp. 132-135. (In Japanese, with English summary).
- 37) Negoro, K. (1968) : An analytical study of diatom shells in the bottom deposits of Lake Yogo-ko. Mem. Fac. Sci. Kyoto Univ., Ser. Biol., Vol. I, No.2, pp. 121-124.
- 38) Negoro, K. (1968) : Diatoms from some inlandwaters of the Antarctica. Mem. Fac. Sci. Kyoto Univ., Ser. Biol., Vol. □, No. 2, pp. 125-138.
- 39) Negoro, K. (1968) : Second report of the regular limnological survey of Lake Biwa (1967). I. Plankton in general and phytoplankton. Mem. Fac. Sci. Kyoto Univ., Ser. Biol., Vol. □, No. 1, pp. 72-91.
- 40) Negoro, K. (1968) : The Phytoplankton of Lake Biwa. S. Kitamura's "Flora Ohmiensis", pp. 275-330, 18 pls. and 177 figs. (In Japanese).
- 41) Negoro, K. (1969) : The diatom-vegetation of the River Yoshino-gawa, Nara Prefecture. Mem. Fac. Sci. Kyoto Univ., Ser. Biol., Vol. □, No. 2, pp. 184-193
- 42) Negoro, K. (1974) : The algal vegetation of some rivers in the Ryu-kyu (Loo-choo) Islands and Formosa. Ecol. Stud. Nat. Cons. Ryukyu Isl., Vol. I, pp. 145-159. (In Japanese, with English summary).
- 43) Negoro, K. (1974) : Eutrophication and plankton in Lake Biwa. T. Fujinaga's "The Dynamic Phase of Lake Biwa", pp. 181-199. (In Japanese).
- 44) Negoro, K. and Masayuki Watanabe (1977) : On the phytoplankton of Lake Akan, Hokkaido. Sorui (Bull. Jap. Soc. Phycol.), Vol. **25**, pp. 221-237. (In Japanese, with English summary).
- 45) Tomoda, Yoshio and K. Negoro (1979) : The newly obtained, paleobiological materials from the Kobiwako Group (Tertiary and quaternary Deposits) of Lake Biwa . Chikyu-kagaku (Earth Science), Vol. **33**, pp. 236-240. (In Japanese).

- 46) Negoro, K. (1979) : Studies on the relation between the water pollution and the algal flora in Lake Biwa and its effluent the River Yodo, with special reference to the diatom flora as an indicator of the pollution (The First Report). Report of the Environmental Pollution Research Institute, Kinki University, No. **7**, pp. 15-20. (In Japanese, with English summary).
- 47) Negoro, K. (1980) : Ditto (The Second Report). Rep. Envir. Poll. Res. Inst., Kinki Univ., No. **8**, pp. 5-11. (In Japanese, with English summary).
- 48) Negoro, K. (1981) : Ditto (The Third Report). Rep. Envir. Poll. Res. Inst., Kinki Univ., No. **9**, pp. 19-22. (In Japanese, with English summary).
- 49) Negoro, K. (1981) : The diatom flora of a pond (so-called "FarmPond") in the experimental farm of the Kinki University, at Yuasa, Wakayama Prefecture, Japan. Mem. Fac. Agr. Kinki Univ., Vol. **14**, pp. 1-42.
- 50) Negoro, K. (1981) : Fossil diatoms of the Kobiwako Group viz. Ancient Deposits of Lake Biwa. Acta Phytotax. Geobot., Vol. **32**, pp. 90-104.
- 51) Negoro, K. (1981) : Fossil diatoms from the Katana Formation of the Kobiwako Group. Acta Phytotax. Geobot., Vol. **32**, pp. 183-191. (In Japanese, with English summary).
- 52) Negoro, K. (1981) : Studies on the phytoplankton of Lake Biwa. Verh. Internat. Verein. Limnol., Bd. **21**, S. 574-583.
- 53) Negoro, K. (1981) : On the disorder of plankton calendar (plankton periodicity) in Lake Biwa, followed by the increased eutrophication. Water-Temperature Research, Vol. **25**, pp. 15-19. (In Japanese).
- 54) Negoro, K. and Toshikazu Gotoh (1981) : Fossil diatoms of the Agé Group, one of the Pliocene freshwater deposits in Japan. (The First Report). Bulletin Mizunami Fossil Museum, No. **8**, pp. 77-103. (In Japanese, with English summary).
- 55) Negoro, K. und T. Gotoh (1983) : *Gyrosigma exoticumoides* sp. nov., eine neue brackische Diatomee aus Japan. Acta Phytotax. Geobot., Vol. **34**, pp. 87-90.
- 56) Negoro, K. and T. Gotoh (1983) : Two new diatoms of the Genus *Navicula* from the River Yura, Kyoto Prefecture, Japan. Acta Phytotax. Geobot., Vol. **34**, pp. 91-93.
- 57) Negoro, K. and T. Gotoh (1983) : Diatom vegetation of the River Yura. Mem. Fac. Agr. Kinki Univ., Vol. **16**, pp. 67-118. (In Japanese, with English summary).
- 58) Gotoh, T. and K. Negoro, (1986) : Diatom vegetation of the less polluted river, the U-kawa River, Kyoto Prefecture. Jap. Jour. Limnol., Vol. **47**, pp. 77-86. (In Japanese, with English summary).
- 59) Gotoh, T. and K. Negoro (1986) : Diatom vegetation of the Totsukawa, Kumano-gawa River System. Jap. Jour. Limnol., Vol. **47**, pp. 143-153. (In Japanese, with English summary).
- 60) Negoro, K. and Masako Higashino (1986) : Diatom vegetation of paddy fields in Japan.

Report I. Diatom vegetation of paddy fields in the vicinity of Sakurai City, Nara Prefecture. Diatom, Vol. **2**, pp. 1-8.

- 61) Negoro, K. (1986) : *Melosira hustedti* Krasske from Lake Biwa, as one of the subfossil diatoms in the bottom deposits. Diatom, Vol. **2**, pp. 19-21.
- 62) Negoro, K. (1986) : Dr. Friedrich Hustedt, Famous Excellent Diatomologist. His life and work. Diatom, Vol. **2**, pp. 187-190. (In Japanese).
- 63) The southern, tropical elements of the diatom-flora of Japan. Acta Phytotax. Geobot., Vol. **38**, pp. 7-15. (In Japanese, with English summary).